

Mpox

Information for Blood Programs



Can Mpox be transmitted via blood transfusion?

There is currently no evidence to suggest that Mpox can be spread through blood transfusion. Donor selection criteria excludes donors who are unwell or exhibiting symptoms.



Can you donate blood after recovering from a Mpox infection?

Infected donors should be deferred until all symptoms are fully resolved, and a minimum of 21 days after the onset of symptoms.

What about close contacts who are asymptomatic?

As a precaution, high-risk close contacts should not donate blood for a minimum of 21 days from their last contact with the known case or contaminated material.



Is there a vaccination deferral period?

Individuals who have recently received a vaccine for Mpox may be temporarily deferred from donating - the deferral timeframe will depend on the type of vaccine received.

Information current as at September 2022

For more information refer to the following resources:

- WHO - [Monkeypox Q&A](#)
- WHO - [Multi-country monkeypox outbreak in non-endemic countries](#)
- FDA - [Information for blood establishments regarding the monkeypox virus and blood donation](#)

All information shared by GAP is provided "as is" without any warranty of fitness for any particular purpose. The advice, guidance and tools provided by GAP are based on GAP's general knowledge and expertise; they are not developed in light of a particular legal regime or national context. Reliance on the information is at the user's sole risk and GAP expressly disclaims any liability of any nature arising out of the use of information. All copyright and intellectual property in the information remains vested in GAP or the member organisation that provided it, as applicable.